

LET'S TALK ABOUT HOW YOU CURRENTLY MANAGE MIGRAINE ATTACKS

UBRELVY
(ubrogepant) tablets 50mg/100mg

Fill out the questions below and bring your answers to your next appointment.

How many migraine attacks do you have each month?



1



2



3



4



5



6



7+

What treatments—both prescription and over the counter—do you currently take for your migraine attacks when they strike?

Do you ever take multiple types of medication to treat a migraine attack?

☐ Yes

☐ No

ASK YOUR HEALTHCARE PROVIDER: Can I take UBRELVY with other migraine medications?

Which symptoms do you experience during a migraine attack?



Head Pain



Light Sensitivity



Sound Sensitivity



Nausea



All of these

ASK YOUR HEALTHCARE PROVIDER: Does UBRELVY treat migraine pain and symptoms?

How often do you question taking your migraine medication because you're in a place where you feel you can't take it?

Never

For a few of my migraine attacks

For many of my migraine attacks

For all of my migraine attacks

ASK YOUR HEALTHCARE PROVIDER: Are there any side effects with UBRELVY?

Do you ever wait to see if your migraine will be "bad enough" before treating it with prescription medication?
If so, do you still get the relief you need?

Never

For a few of my migraine attacks

For many of my migraine attacks

For all of my migraine attacks

ASK YOUR HEALTHCARE PROVIDER: If I don't take my medication right away, will UBRELVY still work?

☒ **TAKE THIS WITH YOU AND ASK YOUR HEALTHCARE PROVIDER ABOUT UBRELVY TODAY!**

What is UBRELVY® (ubrogepant)?

UBRELVY is a prescription medicine used for the acute treatment of migraine attacks with or without aura in adults. UBRELVY is not used to prevent migraine headaches.

IMPORTANT SAFETY INFORMATION

Do not take UBRELVY if you are taking medicines known as strong CYP3A4 inhibitors, such as ketoconazole, clarithromycin, or itraconazole, or if you are allergic to UBRELVY or any of its ingredients.

Before taking UBRELVY, tell your healthcare provider about all your medical conditions, including if you:

- Have high blood pressure
- Have circulation problems in your fingers and toes
- Have liver problems
- Have kidney problems
- Are pregnant or plan to become pregnant
- Are breastfeeding or plan to breastfeed

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Your healthcare provider can tell you if it is safe to take UBRELVY with other medicines.

UBRELVY may cause serious side effects, including:

- **Allergic reactions:** Most reactions happened within hours after taking UBRELVY and were not serious. Some reactions may occur days after taking UBRELVY. Call your healthcare provider or get emergency help right away if you have swelling of the face, mouth, tongue, or throat or trouble breathing.
- **High blood pressure:** New or worsening of high blood pressure can happen. Contact your healthcare provider if you have an increase in blood pressure.
- **Raynaud's phenomenon:** A type of circulation problem can worsen or happen. Raynaud's phenomenon can lead to your fingers or toes feeling numb, cool, or painful, or changing color from pale, to blue, to red. Contact your healthcare provider if these symptoms occur.

The most common side effects of UBRELVY are nausea (4%) and sleepiness (3%). These are not all of the possible side effects of UBRELVY.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/PatientAccessSupport to learn more.

UBRELVY® and its design are registered trademarks of Allergan Pharmaceuticals International Limited, an AbbVie company.
© 2025 AbbVie. All rights reserved. US-UBR-250190